



HPV Prevention
and Control Board



**HOSPITAL
UNIVERSITARIO
SAN IGNACIO**

CIENCIA Y TECNOLOGIA CON PROYECCION SOCIAL


Women's retention and follow-up in cervical cancer screening: a Colombian experience

Raúl Murillo


May 2026

No conflicts of interest to declare

Lost to follow-up (diagnostic workup and treatment) among women with HSIL (1)



5-year follow-up compliance after call and recall in the ESTAMPA study



Follow-up compliance among HPV-positive women in the implementation of HPV self-collection within a screen and treat algorithm

A community trial: four municipalities in two regions



Population
Invited: 1400
Attendant: 532

Compliance	Self-collection		Clinician-collection		<i>p-value</i>
	%	95% CI	%	95% CI	
Triage/treatment visit (2 weeks)					
Total	86,7	78.4-92.7	97,2	85.5-99.9	0,079
Delivery strategy					
Campaign	84,1	72.7-92.1	97,1	85.1-99.9	0,050
Routine screening	91,4	76.9-98.2	100.0	---	0,760
Follow-up visit (6 months)					
Total	71,4	61.4-80.1	69,4	51.9-83.7	0.823
Delivery strategy					
Campaign	76,2	63.8-86.0	68,6	50.7-83.1	0.413
Routine screening	62.9	44.9-78.5	100.0	---	0,446

1-year follow-up compliance (Putumayo): 42,5%

Key messages

- The lost to follow-up for diagnostic workup and treatment about 30%
- The lost to follow-up for a second round of screening over 50%
- The main barriers are related to health services
- The main determinants are the number of visits and the elapsed time between visits
- Different realities require different approaches



Thank you!
